
MRI Fusion Biopsy Instructions

YOUR PHYSICIAN HAS ORDERED AN MRI FUSION BIOPSY OF THE PROSTATE

PRIOR TO YOUR MRI FUSION BIOPSY

- **DO NOT TAKE BLOOD THINNERS SUCH AS:** Aspirin, Aleve, Ibuprofen, Motrin, Plavix, Coumadin or any other blood thinners for 7 days prior to your procedure.
If a pain reliever is needed, 1000mg of acetaminophen (Tylenol) may be taken.
- Do not consume alcohol for 72 hours prior to the biopsy.
- Abstain from sexual activity for 72 hours prior to the biopsy.

DAY OF THE MRI FUSION BIOPSY

@UroPartners Surgery Center 2750 S. River Rd., Des Plaines, IL 60018

DATE/DAY: _____

ARRIVAL TIME: Given to you the afternoon before your procedure.

DO NOT EAT OR DRINK ANYTHING STARTING AT MIDNIGHT PRIOR TO YOUR SCHEDULED BIOPSY.

- Your rectum must be emptied for your procedure. **In order to do so, please use a fleet enema; use this fleet enema two hours before bedtime.**
- Please take the antibiotic Levaquin (unless allergic) by mouth that your physician prescribed for you one to two hours before your procedure.
- Anesthesia will be administered; **you must have a responsible adult over the age of 18 to drive you home from your procedure.**

If you have any questions regarding your procedure preparations or medications, please call our center at 224-612-7000.

ADDRESS OF THE CENTER

**UroPartners Surgery Center
2750 S. River Rd.
Des Plaines, IL 60018**